

Arp2/3 Complex Regulation

Antibody Sampler Kit

Catalog # AK6390

Kit Components:

Catalog#	Description	Host	Size	Applications	Species Reactivity	MW (kDa)
AP3861	Arp2 (C-terminal region)	Rabbit pAb	50 μ l	WB, E, ICC	H, R, M, C, F	44
AP3871	Arp2 (Thr-237/Thr-238), phospho-specific	Rabbit pAb	50 μ l	WB, E, ICC	H, R, M, C, F	44
WP2101	WASP / N-WASP	Rabbit pAb	50 μ l	WB, E, ICC	H, R, M	65
WP2601	N-WASP (Tyr-256), phospho-specific [Conserved site]	Rabbit pAb	50 μ l	WB, E, ICC	H, R, M	65
CP2581	Coronin-1B (C-terminus)	Rabbit pAb	50 μ l	WB, E, ICC	H, R, M	60
CP2621	Coronin-1B (Ser-2), phospho-specific	Rabbit pAb	50 μ l	WB, E, ICC	H, R, M	60

Applications: WB = western blot, E = ELISA, ICC = immunocytochemistry. Species: H = Human, R = Rat, M = Mouse, C = Chicken, F = Fish

Kit Summary:

The Arp2/3 complex regulation antibody sampler kit can be used to examine phosphorylation of the Arp2/3 complex subunit Arp2, as well as the Arp2/3 complex regulators N-WASP/WASP and Coronin-1B. The kit contains antibodies to monitor total expression levels of each of these proteins relative to site-specific phosphorylation.

Buffers and Storage:

Rabbit polyclonal antibodies are each supplied in 50 μ l phosphate-buffered saline, 50% glycerol, 1 mg/ml BSA, and 0.05% sodium azide.

Store at -20°C . Do not aliquot. Stable for 1 year.

Background:


Cellular morphology, adhesion, and motility occur through dynamic reorganization of actin-based superstructures. Actin-binding proteins are critical for regulating actin polymerization and superstructure formation. The Arp2/3 complex is an actin polymerization-inducing complex that includes Arp2, Arp3, p41-Arc, p34-Arc, p21-Arc, p20-Arc, and p16-Arc. Several nucleation promoting factors, such as WASPs and coronins, regulate the activity of the Arp2/3 complex. WASP is tyrosine phosphorylated at tyrosine 291 after antigen receptor activation in B-cells and collagen stimulation of platelets. Phosphorylation of the analogous site in N-WASP (Tyr-256) stimulates its activity, reduces nuclear N-WASP, and is required for neurite extension. Both Coronin-1B and Coronin-1A interaction with Arp2/3 complex may be regulated by phosphorylation. PKC phosphorylates the N-terminus at Ser-2, and this phosphorylation reduces interactions with Arp2/3 leading to diminished cell motility. The Arp2/3 complex is also regulated by phosphorylation. Arp2 has two phosphosites, Thr-237 and Thr-238, which are evolutionarily conserved. These sites are phosphorylated along with Tyr-202 in response to growth factor stimulation. These phosphorylation events may regulate binding to the pointed end of actin filaments, and alanine substitutions of these Arp2 phosphosites inhibit membrane protrusions.

References:

Wu, X. et al. (2004) J Biol Chem 279(10):9565.
 Foger, N. et al. (2006) Science 313:839.
 Cai, L. et al. (2007) Cell 128:915.
 LeClaire, L.L. et al (2008). J Cell Biol. 182(4):647.
 Soderling, S.H. (2009). Sci Signal. 2(55):pe5 (Review).

Product References (WP2601) :

Pichot, C.S. et al. (2009) British J. Cancer 1 –10. (WB: Breast Cancer Cell lines)



FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

www.ecmbiosciences.com
 telephone: 859-879-2075
 toll-free: 1-800-859-8202
 info@ecmbiosciences.com

ECMBiosciences