

A431 + EGF (5 min) Lysate

Cat. # AL9301

Size 100 μ l

Lysate Preparation:

A431 cells express approximately 10^6 epidermal growth factor (EGF) receptors at the cell surface. Upon stimulation with EGF, A431 cells exhibit a dramatic increase in phosphorylation of EGF receptors followed by activation of major cell signaling pathways, such as PKB/Akt and MAP kinase pathways. Downstream of these cell signaling pathways, a variety of cytoskeletal, cytoplasmic, and nuclear proteins become phosphorylated at different time points after EGF stimulation.

Confluent cultures of A431 cells were serum starved overnight. Cells were then either left untreated (Cat.# AL9201) or treated with human EGF (100 ng/ml) for 30 minutes at 37°C (Cat.# AL9301). Cells were lysed in 1% SDS, 1.0 mM sodium ortho-vanadate, 10 mM Tris (pH 7.4) buffer. Protein concentration was determined using the BCA method (Pierce) before diluting to final concentration and buffer.

Applications:


WB 20 μ l/lane

End user should determine optimal quantity for their particular applications and experiments.

Related Products:

AL9201 A431 EGF Ctrl Lysate
AL9001 A431 Calyculin A Ctrl Lysate
AL9101 A431 + Calyculin A (30min) Lysate
AL9401 A431 Pervanadate Ctrl Lysate
AL9501 A431 + Pervanadate Lysate

Application Data:



Western blot analysis of A431 cells (20 μ g/lane) serum starved overnight and treated with EGF (100 ng/ml) for 5 min. The blot was probed with anti-paxillin (PM1071) or anti-paxillin (Ser-178).

Buffer and Storage:

Cell Lysates are supplied at a concentration of 1 mg/ml in electrophoresis sample buffer (62.5 mM Tris pH 6.8, 2% SDS, 5% glycerol, 0.003% bromophenol blue, 0.9% β -mercaptoethanol). Store at -20°C . Do not boil or dilute. Stable for 1 year.

FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

www.ecmbiosciences.com
telephone: 859-879-2075
toll-free: 1-800-859-8202
tech: info@ecmbiosciences.com

ECMBiosciences

Rev 10/19/2007