

ATM (Ser-794), phospho-specific

Cat. # AP3631

Host Rabbit Polyclonal


Size 100 µl

Background:

Ataxia telangiectasia mutated kinase (ATM) is a serine/threonine kinase that regulates cell cycle checkpoints and DNA repair. Mutations of ATM cause a spectrum of defects ranging from neurodegeneration to cancer predisposition. Activation of ATM after DNA damage involves Cdk5 mediated phosphorylation of Ser-794 followed by autophosphorylation at Ser-1891. Active ATM kinase regulates a number of proteins involved in cell cycle checkpoint control, apoptosis and DNA repair. The Cdk5-ATM pathway regulates phosphorylation and function of the ATM targets p53 and H2AX in postmitotic neurons. Other known substrates of ATM include Chk2, Chk1, CtIP, 4E-BP1, BRCA1, RPA3, SMC1, FANCD2, Rad17, Artemis, Nbs1, and the I-2 regulatory subunit of PP1. Thus, activation of Cdk5 by DNA damage may be an important initiator of ATM-dependent regulation of cell cycle checkpoints.

References

- Shiloh, Y. (1997) *Annu Rev Genet.* 31:635.
 Lee, J.H. & Paull, T.T. (2007) *Oncogene* 26:7741.
 Tian, B. et al. (2009) *Nat Cell Biol.* 11:211.



FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

web: www.ecmbiosciences.com

telephone: 859-879-2075

email: info@ecmbiosciences.com

toll-free: 1-800-859-8202



ATM (Ser-794), phospho-specific

Cat. # AP3631

Host Rabbit Polyclonal

Size 100 µl

Immunogen:

Phospho-ATM (Ser-794) synthetic peptide (coupled to carrier protein) corresponding to amino acids surrounding Ser-794 in human ATM. This sequence is well conserved in rat and mouse ATM.

Buffer and Storage:

Rabbit polyclonal, affinity-purified antibody is supplied in 100µl phosphate-buffered saline, 50% glycerol, 1 mg/ml BSA, and 0.05% sodium azide. Store at -20°C. Do not aliquot. Stable for 1 year.

Applications:

WB 1:1000

ELISA 1:2000

ICC 1:200

End user should determine optimal dilution for their particular applications and experiments.
Western blot membranes were incubated with diluted antibody in 5% non-fat milk, PBS, 0.04% Tween20 for 1hour at room temperature.

Specificity:

This antibody was affinity purified using phospho-ATM (Ser-794) peptide (without carrier). The antibody detects a 370 kDa* band corresponding to ATM on SDS-PAGE immunoblots of calyculin A treated Jurkat, A431, HeLa, and rat PC12 cells. This reactivity is removed after lambda phosphatase treatment.

*All molecular weights (MW) are confirmed by comparison to Bio-Rad Rainbow Markers and to western blot mobilities of known proteins with similar MW.

Related Products:

- AM3611 ATM (C-terminal region) Mouse Monoclonal
- AM3661 ATM (Ser-1981), phospho-specific Mouse Monoclonal
- CM2361 Cdk5 Mouse Monoclonal
- CM2311 Cdk1 (Tyr-15), phospho-specific [Conserved site] Mouse Monoclonal
- MS3001 Donkey Anti-Mouse Ig:HRP
- RS3251 Mouse Anti-Rabbit Ig Light-Chain Specific:HRP Mouse Monoclonal

FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

web: www.ecmbiosciences.com

telephone: 859-879-2075

email: info@ecmbiosciences.com

toll-free: 1-800-859-8202

ECMBiosciences