

Cdk1 (N-terminal region)

Cat. # CM2261

Host Mouse Monoclonal IgG1

Size 100 µl

Background:

Cyclin-dependent kinases (Cdks) are a family of serine/threonine kinases that require association with regulatory subunits known as cyclins for activation. In addition, post-translational phosphorylation and dephosphorylation events regulate Cdk activity. Phosphorylation of Thr-160 in the T loop by Cdk-activating kinase (CAK) is an obligatory step in kinase activation. By contrast, phosphorylation of the Thr-14 and Tyr-15 residues by the Wee1 family of dual specificity kinases is inhibitory for the Cdks, and dephosphorylation of these residues by the Cdc25 family of phosphatases coincides with Cdk activation. Alternatively, Cdk5 appears to require different mechanisms for activation. This Cdk is activated through association with specific activators, including p35, p39, and p67. Cdk5 is primarily activated in neuronal cells, and only c-Abl kinase, rather than Wee family members, have been shown to phosphorylate Tyr-15 to regulate its activity.

References

- Poon, R.Y.C. et al. (1997) J Biol. Chem. 272(9):5703.
Zukerberg, L. R. (2000) Neuron 26:633.
Lee, J.H. et al. (2008) J Biol. Chem. May 19 epub.

Immunogen:

This antibody was generated from a recombinant human Cdk1 protein that included amino acids residues in the N-terminal region. This sequence is conserved in human and rat Cdk1, and has low homology to other Cdk family members.


Applications:

WB	1:1000
ELISA	1:2000
ICC	1:100

End user should determine optimal dilution for their particular applications and experiments. Western blot membranes were incubated with diluted antibody in 5% non-fat milk, PBS, 0.04% Tween20 for 1hour at room temperature.

Related Products:

- CM2311 Cdk1 (Tyr-15), phospho-specific [Conserved site] Mouse Monoclonal
CM2361 Cdk5 Mouse Monoclonal



Western blot analysis of human SYF fibroblasts before (lanes 1 & 3) and after (lanes 2 & 4) treatment with alkaline phosphatase. The blots were probed with anti-Cdk1 (N-terminal region) antibody (lanes 1 & 2) or anti-Cdk1 (Tyr-15) phospho-specific antibody (lanes 3 & 4).

Buffer and Storage:

Mouse monoclonal antibody purified with protein A chromatography is supplied in 100µl phosphate-buffered saline, 50% glycerol, 1 mg/ml BSA, and 0.05% sodium azide. Store at -20°C. Do not aliquot. Stable for 1 year.

Specificity:

The antibody detects a 34 kDa* band corresponding to Cdk1 on SDS-PAGE immunoblots of human SYF and HeLa cells. The antibody does not detect other Cdk family members, such as Cdk2 and Cdk5.

*All molecular weights (MW) are confirmed by comparison to Bio-Rad Rainbow Markers and to western blot mobilities of known proteins with similar MW.

FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

www.ecmbiosciences.com
telephone: 859-879-2075
toll-free: 1-800-859-8202
tech: info@ecmbiosciences.com

ECMBiosciences

Rev 7/14/2008