

Caveolin-1 (N-terminal region)

Cat. # CP2781

Host Rabbit Polyclonal


Size 100 µl

Background:


Caveolins are the primary structural components of the plasma membrane microdomains, caveolae. Three members of the caveolin family (caveolin-1, -2, and -3) have been identified, and each has distinct expression patterns. Caveolins are involved in diverse biological functions, including vesicular trafficking, cholesterol homeostasis, cell adhesion and apoptosis. Caveolins can interact with various signaling molecules, including G-proteins, receptor tyrosine kinases, PKCs, and Src family kinases. Phosphorylation at Tyr-14 is essential for caveolin association with SH2 or PTB domain-containing adaptor proteins, while phosphorylation at Ser-80 regulates caveolin binding to the ER membrane and entry into the secretory pathway.

References

- Okamoto, T. et al. (1998) J. Biol. Chem. 273:5419.
Smart, E. J. et al. (1999) Mol. Cell. Biol. 19:7289.
Nomura, R. et al. (1999) Mol. Biol. Cell 10:975.
Schlegel, A. et al. (2001) J. Biol. Chem. 276:4398.



Immunocytochemical labeling of caveolin-1 in paraformaldehyde-fixed and NP-40-permeabilized rabbit spleen fibroblasts. The cells were labeled with rabbit polyclonal Caveolin-1 (N-terminal region) and mouse monoclonal Caveolin-1 antibodies, and detected using appropriate secondary antibodies conjugated to Cy3. Phase contrast images (left) and immunofluorescent images (right).



Western blot image of human A431 cells unstimulated (lanes 1, 3, & 5) or stimulated with pervanadate (1 mM) for 30 min (lanes 2, 4, & 6). The blots were probed with rabbit polyclonal caveolin-1 (N-term.) (lanes 1 & 2), mouse monoclonal caveolin-1 (Tyr-14) (lanes 3 & 4) or mouse monoclonal caveolin-1 (lanes 5 & 6).

FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

www.ecmbiosciences.com
telephone: 859-879-2075
toll-free: 1-800-859-8202
email: info@ecmbiosciences.com

ECMBiosciences

Rev 3/3/2009

Caveolin-1 (N-terminal region)

Cat. # CP2781

Host Rabbit Polyclonal

Size 100 µl

Immunogen:

The antibody was generated from a recombinant protein that includes amino acids in the N-terminal region of human Caveolin-1.

Buffer and Storage:

Rabbit polyclonal, affinity-purified antibody is supplied in 100 µl phosphate-buffered saline, 50% glycerol, 1 mg/ml BSA, and 0.05% sodium azide. Store at -20°C. Do not aliquot. Stable for 1 year.

Applications:

WB 1:2000

ELISA 1:4000

ICC 1:100

End user should determine optimal dilution for their particular applications and experiments.
Western blot membranes were incubated with diluted antibody in 5% non-fat milk, PBS, 0.04% Tween20 for 1 hour at room temperature.

Specificity:

This antibody detects a 21 kDa* protein corresponding to the apparent molecular mass of caveolin-1 on SDS-PAGE immunoblots of human A431 cells and rabbit spleen fibroblasts.

*All molecular weights (MW) are confirmed by comparison to Bio-Rad Rainbow Markers and to western blot mobilities of known proteins with similar MW.

Related Products:

- CM2811 Caveolin-1 Mouse Monoclonal
- CM2831 Caveolin-1 (Tyr-14), phospho-specific Mouse Monoclonal
- EP1911 EGFR (Ser-967), phospho-specific Rabbit Polyclonal
- EM1991 EGFR (Tyr-1101), phospho-specific Mouse Monoclonal
- EP1931 EGFR (Ser-1142), phospho-specific Rabbit Polyclonal
- SM2611 c-Src (Tyr-530), phospho-specific [Conserved site] Mouse

FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

www.ecmbiosciences.com
telephone: 859-879-2075
toll-free: 1-800-859-8202
email: info@ecmbiosciences.com

ECMBiosciences

Rev 3/3/2009