

β -Catenin (C-terminal) Blocking Peptide

Cat. # **CX1205**


Size **50 μ g**

Background:

β -Catenin is a 92 kDa protein that binds to the cytoplasmic tail of E-Cadherin. The cadherins, transmembrane adhesion molecules, are found with catenins at adherens junctions. Deletions in the cytoplasmic domain of E-Cadherin eliminate catenin binding and result in a loss of cell adhesion. Tyrosine phosphorylation of β -Catenin can regulate its interaction with critical components of adherens junctions. Both Fer and Fyn kinases phosphorylate tyrosine 142 *in vitro*. Overexpression of these kinases in epithelial cells disrupts interactions between α - and β -Catenins. The phosphorylation of tyrosine 142 may act as a switch from the transcriptional to the adhesive role of β -Catenin. Src family kinases can also phosphorylate tyrosine 86 and 654 in β -Catenin. The Tyr-654 phosphorylation regulates β -Catenin binding to E-cadherin. Thus, site-specific tyrosine phosphorylation of β -Catenin may regulate protein-protein interactions, leading to changes in cell adhesion.

References:

Ozawa, M. et al. (1990) Proc. Natl. Acad. Sci. USA 87:4246.
Roura, S. et al. (1999) J Biol Chem. 274(51):36734.
Piedra, J. et al. (2003) Mol. Cell. Biol. 23(7):2287-2297.
Brembeck, F.H. et al. (2004) Genes Dev. 18(18):2225-2230.



Peptide Sequence:

A synthetic peptide corresponding to amino acid residues in the C-terminal region of human β -Catenin. This peptide sequence has homology to the C-terminal region of γ -Catenin and is highly conserved in rat and mouse β - and γ -Catenins.

Buffer and Storage:

Blocking Peptide is supplied in 50 μ l phosphate-buffered saline and 0.05% sodium azide. Store at -20° C. Stable for 1 year.

Applications:

Antibody Blocking 1 μ g/ml¹
ELISA 10-100 ng/well²

End user should determine optimal concentration dependent on the concentration of the antibody.

¹Recommended for blocking antibody reactivity in Western blot and immunocytochemistry.

²Established in 96-well Nunc immunoplates where peptide was bound to plates for 2 hrs in 0.1 M sodium carbonate buffer, pH 8.5.

Specificity:

The peptide is specifically recognized by anti- β -Catenin (C-terminal) antibody (CP1201) in ELISA, and has been shown to block the reactivity of CP1201 during Western blot. In addition, the peptide is recommended for use in blocking CP1201 reactivity in immunocytochemistry.

Related Products:

CP1201 β -Catenin (C-terminal) Rabbit Polyclonal
CP1061 β -Catenin (N-terminal) Rabbit Polyclonal

CP1081 β -Catenin (Tyr-142) (γ -Catenin (Tyr-133)), phospho-specific Rabbit Polyclonal
CP1191 β -Catenin (Tyr-86), phospho-specific Rabbit Polyclonal

FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

www.ecmbiosciences.com
telephone: 859-879-2075
toll-free: 1-800-859-8202
tech: info@ecmbiosciences.com

ECMBiosciences

Rev 10/06