

Rabbit Fibroblast Calyculin A Control Lysate

Cat. # FL9671

Size 100 μ l

Lysate Preparation:

Calyculin A is a serine/threonine phosphatase inhibitor that inhibits the activity of protein phosphatases PP1 and PP2A. Cells treated with calyculin A for 30 minutes can undergo significant threonine phosphorylation, as shown by western blotting using anti-Phosphoserine/threonine, cat.# PP2551, as compared to untreated, control cell lysates.

Fibroblast from rabbit spleen were harvested to generate a primary fibroblast cell line. Confluent cultures of rabbit fibroblasts were serum starved for 2 hours. Cells were then either left untreated (Cat.# FL9671) or treated with Calyculin A (100 nM) for 30 minutes at 37°C (cat.# FL9681). Cells were lysed in 1% SDS, 1.0 mM sodium ortho-vanadate, 1 mM sodium fluoride in 10 mM Tris (pH 7.4) buffer. Protein concentration was determined using the BCA method (Pierce) before diluting to final concentration and buffer.

Buffer and Storage:

Cell Lysates are supplied at a concentration of 1 mg/ml in electrophoresis sample buffer (62.5 mM Tris pH 6.8, 2% SDS, 5% glycerol, 0.003% bromophenol blue, 0.9% β -mercaptoethanol). Store at -20°C . Do not boil or dilute. Stable for 1 year.

Applications:

WB 20 μ l/lane

End user should determine optimal quantity for their particular applications and experiments.

Related Products:

FL9681 Rabbit Fibroblast + Calyculin A Lysate
FL9691 Rabbit Fibroblast Pervanadate Control Lysate
FL9701 Rabbit Fibroblast + Pervanadate Lysate
PK6330 Phospho-Tyrosine, Serine, Threonine Antibody Sampler Kit
PM3801 Anti-Phosphoserine/threonine Mouse Monoclonal
PP2551 Anti-Phosphoserine/threonine Rabbit Polyclonal

FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

www.ecmbiosciences.com
telephone: 859-879-2075
toll-free: 1-800-859-8202
email: info@ecmbiosciences.com

ECMBiosciences

Rev 6/8/2010