

Jurkat Camptothecin Control Lysate

Cat. # JL9531

Size 100 μ l

Lysate Preparation:

The camptothecin treatment of human Jurkat T-Cell leukemia cells can be used as a western blot positive control for detecting antigens expressed during apoptosis. The treated cells undergo apoptosis as determined by identification of specific apoptotic cleavage products, such as PARP and caspase-3 cleavage.

Confluent cultures of Jurkat cells were either left untreated (Cat.# JL9531) or treated with Camptothecin (4 μ M) for 4 hours at 37°C (cat.# JL9541). Cells were lysed in 1% SDS, 1.0 mM sodium ortho-vanadate, 1 mM sodium fluoride in 10 mM Tris (pH 7.4) buffer. Protein concentration was determined using the BCA method (Pierce) before diluting to final concentration and buffer.

Buffer and Storage:

Cell Lysates are supplied at a concentration of 1 mg/ml in electrophoresis sample buffer (62.5 mM Tris pH 6.8, 2% SDS, 5% glycerol, 0.003% bromophenol blue, 0.9% β -mercaptoethanol). Store at -20° C. Do not boil or dilute. Stable for 1 year.

Applications:

WB 20 μ l/lane

End user should determine optimal quantity for their particular applications and experiments.

Related Products:

JL9541 Jurkat Camptothecin-induced Apoptosis Lysate
JL9251 Jurkat Calyculin A Control Lysate
JL9261 Jurkat + Calyculin A Lysate
JL9401 Jurkat Pervanadate Control Lysate
JL9501 Jurkat + Pervanadate Lysate
CK6360 Caspase Family Antibody Sampler Kit

FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

www.ecmbiosciences.com
telephone: 859-879-2075
toll-free: 1-800-859-8202
email: info@ecmbiosciences.com

ECMBiosciences

Rev 6/8/2010