

p38 MAP Kinase (Thr-180/Tyr-182), phospho-specific

Cat. # PM1391

Host Mouse Monoclonal IgG1


Size 100 µl

Background:

p38 MAP kinase (MAPK), also called RK, CSBP, and SAPK2a, is the mammalian orthologue of the yeast HOG kinase. This family of kinases participates in signaling cascades that control cellular responses to cytokines and stress. Four isoforms of p38 MAPK (α , β , γ , δ) have been identified. Similar to the SAPK/JNK pathway, p38 MAPK is activated by a variety of cellular stresses including osmotic shock, inflammatory cytokines, lipopolysaccharides, UV light, and growth factors. MKK3 and SEK activate p38 MAPK by dual phosphorylation at threonine 180 and tyrosine 182. Activated p38 MAPK has been shown to phosphorylate and activate MAPKAP kinase 2 and to phosphorylate the transcription factors ATF-2, Max and MEF2.

References

- Han, J. et al. (1994) Science 265:808.
 Lee, J. C. et al. (1994) Nature 372:739.
 Rouse, J. et al. (1994) Cell 78:1027.
 Freshney, N. W. et al. (1994) Cell 78:1039.
 Raingeaud, J. et al. (1995) J. Biol. Chem. 270:7420.



FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

web: www.ecmbiosciences.com

telephone: 859-879-2075

email: info@ecmbiosciences.com

toll-free: 1-800-859-8202



p38 MAP Kinase (Thr-180/Tyr-182), phospho-specific

Cat. #	PM1391
Host	Mouse Monoclonal IgG1
Size	100 μ l

Immunogen:

Clone M139 was generated from a phospho-p38 α (Thr-180/Tyr-182) synthetic peptide (coupled to KLH) corresponding to amino acid residues around threonine 180 and tyrosine 182 of rat p38 α . This peptide sequence is highly conserved in the p38 β , γ , and δ MAPKs, and is identical in human and mouse p38 α .

Buffer and Storage:

Mouse monoclonal purified with protein A chromatography is supplied in 100 μ l phosphate-buffered saline, 50% glycerol, 1 mg/ml BSA, and 0.05% sodium azide. Store at -20°C . Do not aliquot. Stable for 1 year.

Applications:

WB 1:1000
ELISA 1:2000
ICC 1:250

End user should determine optimal dilution for their particular applications and experiments.
Western blot membranes were incubated with diluted antibody in 5% non-fat milk, PBS, 0.04% Tween20 for 1hour at room temperature.

Specificity:

This antibody detects a 42 kDa* protein corresponding to the apparent molecular mass of p38 α on SDS-PAGE immunoblots of pervanadate treated human Jurkat and A431 cells, as well as anisomycin treated human HeLa cells.

*All molecular weights (MW) are confirmed by comparison to Bio-Rad Rainbow Markers and to western blot mobilities of known proteins with similar MW.

Related Products:

PM1381 p38 α MAP Kinase (C-terminal) Mouse Monoclonal
PP3411 p38 α MAP Kinase (Tyr-323), phospho-specific Rabbit Polyclonal
AM1011 Akt (N-terminal Region) Mouse Monoclonal
AM1141 Akt (Ser-473), phospho-specific Mouse Monoclonal
AP1001 Akt (Thr-34), phospho-specific Rabbit Polyclonal
PK6140 p38 MAPK Phospho-Regulation Antibody Sampler Kit

FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

web: www.ecmbiosciences.com

telephone: 859-879-2075

email: info@ecmbiosciences.com

toll-free: 1-800-859-8202

ECMBiosciences