

RhoA (Ser-188), phospho-specific

Cat. # RP1361

Host Rabbit Polyclonal

Size 100 µl

Background:

Rho (A, B, & C) proteins are members of the Ras superfamily of GTPases. These proteins regulate a variety of cellular functions, including cell cycle progression, cytoskeletal rearrangement, and gene expression. Rho cycles between the active GTP-bound form and an inactive GDP-bound form. Interconversion between these forms is controlled by guanine nucleotide exchange factors (GEFs) and GTPase-activating proteins (GAPs). The Rho proteins RhoA, RhoB, and RhoC are highly homologous and contain the consensus amino acid sequences necessary for GDP/GTP-binding and GTPase activity. Post-translational regulation of Rho activity has been shown specifically for RhoA. This Rho protein is phosphorylated *in vitro* on serine 188 by cAMP- and cGMP-dependent kinases (PKA and PKG). Ser-188 phosphorylation enhances RhoGDI binding and inhibits RhoA-mediated stress fiber formation. Thus, Ser-188 is an important site for negative regulation of RhoA activity.

References

- Lang, P. et al. (1996) EMBO J. 15:510.
Dong, J.M. et al. (1998) J. Biol. Chem. 273(35):22554.
Sawada, N. et al. (2001) Biochem. Biophys. Res. Commun. 280:798.
Ellerbroek, S.M. et al. (2003) J. Biol. Chem. 278(21):19023.

Immunogen:

Phospho-RhoA (Ser-188) synthetic peptide (coupled to KLH) corresponding to amino acid residues around serine 188 of human RhoA. This peptide sequence is highly conserved in rat and mouse RhoA, but is not found in RhoB or RhoC proteins.

Applications:

WB 1:1000


ELISA 1:2000

End user should determine optimal dilution for their particular applications and experiments. Western blot membranes were incubated with diluted antibody in 5% non-fat milk, PBS, 0.04% Tween20 for 1 hour at room temperature.

Related Products:

- RP1501 Rho (Central Region) Rabbit Polyclonal
RX1365 phospho-RhoA (Ser-188) Peptide
CM1521 Cdc42 Mouse Monoclonal
RM2721 ROCK-I Mouse Monoclonal
RM2741 ROCK-I (C-terminal), cleavage-specific Mouse Monoclonal
RM2761 ROCK-II (Central region) Mouse Monoclonal

FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



Western blot analysis of human RhoA GST fusion recombinant unphosphorylated (lanes 1 & 3) or phosphorylated with PKA (lanes 2, 4, 5 & 6). The blots were probed with anti-Rho (RP1501; lanes 1 & 2) or with anti-RhoA (Ser-188) (RP1361; lanes 3-6). The latter antibody was used in the presence of no peptide (lanes 3 & 4), phospho-RhoA (Ser-188) peptide (lane 5), or a non-specific phosphoserine peptide (lane 6).

Buffer and Storage:

Rabbit polyclonal, affinity-purified antibody is supplied in 100µl phosphate-buffered saline, 50% glycerol, 1 mg/ml BSA, and 0.05% sodium azide. Store at -20°C. Do not aliquot. Stable for 1 year.

Specificity:

This antibody was affinity purified using phospho-Rho (Ser-188) peptide (without carrier). The antibody detects a RhoA GST fusion protein when phosphorylated with PKA, but does not detect the unphosphorylated recombinant protein.

*All molecular weights (MW) are confirmed by comparison to Bio-Rad Rainbow Markers and to western blot mobilities of known proteins with similar MW.

www.ecmbiosciences.com
telephone: 859-879-2075
toll-free: 1-800-859-8202
email: info@ecmbiosciences.com

ECMBiosciences

Rev 1/2/2008