

SYF c-Src-transformed Control Lysate

Cat. # SL7171

Size 100 μ l

Lysate Preparation:

c-Src was the first proto-oncogenic non-receptor tyrosine kinase characterized in human. The Src family is composed of nine members in vertebrates, including c-Src, Yes, Fgr, Yrk, Fyn, Lyn, Hck, Lck, and Blk. Src-family kinases transduce signals that are involved in the control of a variety of cellular processes, including proliferation, differentiation, motility, and adhesion.

The mouse fibroblast cell line, SYF, is deficient for the c-Src family kinases, c-Src, Yes, and Fyn, while SYF + c-Src overexpress c-Src kinase. This cell line is useful for studies of c-Src mediated phosphorylation when treated with the tyrosine phosphatase inhibitor, pervanadate. Confluent cultures of SYF + c-Src cells were serum starved 2 hrs, and then either left untreated (Cat.# SL7171) or treated with pervanadate (1 mM) for 30 minutes at 37°C (cat.# SL7181). Cells were lysed in 1% SDS, 1.0 mM sodium ortho-vanadate, 10 mM Tris (pH 7.4) buffer. Protein concentration was determined using the BCA method (Pierce) before diluting to final concentration and buffer.

Buffer and Storage:

Cell Lysates are supplied at a concentration of 1 mg/ml in electrophoresis sample buffer (62.5 mM Tris pH 6.8, 2% SDS, 5% glycerol, 0.003% bromophenol blue, 0.9% β -mercaptoethanol). Store at -20°C . Do not boil or dilute. Stable for 1 year.

Applications:

WB 20 μ l/lane

End user should determine optimal quantity for their particular applications and experiments.

Related Products:

SP1371 c-Src (Tyr-215), phospho-specific [Conserved site] Rabbit Polyclonal
SX1375 phospho-c-Src (Tyr-215) Peptide
SL7161 SYF c-Src-transformed Normal Lysate
SL7181 SYF c-Src-transformed + Pervanadate Lysate

FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

www.ecmbiosciences.com
telephone: 859-879-2075
toll-free: 1-800-859-8202
tech: info@ecmbiosciences.com

ECMBiosciences

Rev 10/19/2007