

SHP1 (C-terminal Region)

Cat. # SM1601

Host Mouse Monoclonal IgG1

Size 100 µl

Background:

SHP1 (PTP1C, SH-PTP1, or HCP) is a protein-tyrosine phosphatase (PTP) involved in cell migration, cell proliferation, and immune cell function. This phosphatase contains two N-terminal SH2 domains and a C-terminal phosphatase domain. SHP1 associates with a variety of cytokine and growth factor receptors and regulates signal transduction through dephosphorylation of these receptors or their downstream effectors. Downstream of receptor activation, SHP1 regulates the transcriptional activity stimulated by JAK/Stat and MAPK pathways. SHP1 activity is regulated by both tyrosine and serine phosphorylation. Phosphorylation of Tyr-536 and Tyr-564 stimulates phosphatase activity and promotes interaction with Grb-2. Serine phosphorylation at Ser-591 is mediated by PKC α and leads to inhibition of phosphatase activity. Thus, phosphorylation at tyrosine relative to serine residues may be regulated by different cell signaling pathways to control SHP1 activity.

References

- Jones, M.L. et al. (2004) J. Biol. Chem. 279(39):40475.
Zhang, Z. et al. (2003) J. Biol. Chem. 278(7):4668.
Zhang, J. et al. (2000) Semin. Immunol. 12:361.

Immunogen:

Clone (M160) was generated from a recombinant protein containing amino acids in the C-terminal region of human SHP1. This sequence is highly conserved in rat and mouse SHP1.

Applications:


WB	1:500	ICC	1:100
ELISA	1:2000		
IP	1:100		

End user should determine optimal dilution for their particular applications and experiments. Western blot membranes were incubated with diluted antibody in 5% non-fat milk, PBS, 0.04% Tween20 for 1 hour at room temperature.

Related Products:

SP1571	SHP1 (Tyr-536), phospho-specific	Rabbit Polyclonal
SP1531	SHP1 (Ser-591), phospho-specific	Rabbit Polyclonal
SM1631	SHP2 (N-terminal Region)	Mouse Monoclonal
PP2351	PTP1B (a.a.146-157)	Rabbit Polyclonal
PP2411	PTP1B (Ser-50)	Rabbit Polyclonal
PP2391	PTP1B (Tyr-152)	Rabbit Polyclonal

FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



Western blot analysis of human Jurkat cells treated with pervanadate (1 mM) for 30 min. The blot was exposed to lambda phosphatase (lanes 2 & 4) then probed with anti-SHP1 (C-terminal) antibody (lanes 1 & 2) or anti-SHP1 (Ser-591) antibody (lanes 3-6). The SHP1 (Ser-591) antibody was used in the presence of phospho-SHP1 (Ser-591) peptide (lane 5) or a non-specific phospho-serine peptide (lane 6).

Buffer and Storage:

Mouse monoclonal antibody purified with protein A chromatography is supplied in 100µl phosphate-buffered saline, 50% glycerol, 1 mg/ml BSA, and 0.05% sodium azide. Store at -20°C. Do not aliquot. Stable for 1 year.

Specificity:

The antibody detects a 68 kDa* protein in human A431 and Jurkat cells, and does not cross-react with SHP2.

*All molecular weights (MW) are confirmed by comparison to Bio-Rad Rainbow Markers and to western blot mobilities of known proteins with similar MW.

www.ecmbiosciences.com
telephone: 859-879-2075
toll-free: 1-800-859-8202
tech: info@ecmbiosciences.com

ECMBiosciences

Rev 12/19/2007