

VASP (C-terminal region)

Cat. # VM2771

Host Mouse Monoclonal IgG1


Size 100 µl

Background:


Actin filament tethering and bundling are important mechanisms involved in actin superstructure assembly. The ENA/VASP family includes VASP, mena, and Ena-Vasp-like (EVL). These multidomain proteins localize to the leading edge of filopodia where they associate with AFs, interact with profilin, and compete with capping proteins at the barbed end of AFs. Artificial relocation of VASP from the plasma membrane to mitochondrial membranes inhibits filopodial formation and axon branching, while deletion of all three ENA/VASP proteins produces defects in cortical axon-tract formation. Regulation of VASP protein activity occurs through phosphorylation at Ser-157, Ser-239, and Thr-278. AMPK phosphorylates Thr-278, leading to impaired actin stress fiber assembly and changes in cell morphology.

References

- Krause, M. et al. (2003) *Annu Rev Cell Dev Biol.* 19: 541.
 Blume, C. et al. (2007) *J Biol Chem.* 282(7):4601.
 Applewhite, D.A. et al. (2007). *Mol Biol Cell.* 18(7):2579.



Immunocytochemical labeling of VASP in aldehyde-fixed and NP-40-permeabilized A431 cells. The cells were labeled with mouse monoclonal VASP (C-terminal region) antibody, then the antibody was detected using appropriate secondary antibody conjugated to DyLight 594.



Western blot image of human A431 cells stimulated with calyculin A (100 nM) for 30 min. The blots were untreated (lanes 1 & 3) or treated with lambda phosphatase (lanes 2 & 4), then probed with mouse monoclonal VASP (C-term.) antibody (lanes 1 & 2) or rabbit polyclonal VASP (Thr-278) phospho-specific antibody (lanes 3 & 4).

FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

web: www.ecmbiosciences.com

telephone: 859-879-2075

email: info@ecmbiosciences.com

toll-free: 1-800-859-8202

VASP (C-terminal region)

Cat. # VM2771

Host Mouse Monoclonal IgG1

Size 100 µl

Immunogen:

Clone (M277) was generated from a recombinant protein that includes amino acids from the C-terminal region of human VASP.

Buffer and Storage:

Mouse monoclonal antibody purified with protein A chromatography is supplied in 100µl phosphate-buffered saline, 50% glycerol, 1 mg/ml BSA, and 0.05% sodium azide. Store at -20°C. Do not aliquot. Stable for 1 year.

Applications:

WB 1:1000

ELISA 1:2000

ICC 1:50

End user should determine optimal dilution for their particular applications and experiments.
Western blot membranes were incubated with diluted antibody in 5% non-fat milk, PBS, 0.04% Tween20 for 1hour at room temperature.

Specificity:

This antibody detects 46 and 50 kDa* proteins corresponding to the apparent molecular mass of VASP on SDS-PAGE immunoblots of human A431, HeLa, and HUVEC.

*All molecular weights (MW) are confirmed by comparison to Bio-Rad Rainbow Markers and to western blot mobilities of known proteins with similar MW.

Related Products:

VP2781 VASP (Thr-278), phospho-specific Rabbit Polyclonal

FM2651 Fascin (clone 55K2) Mouse Monoclonal

FP2661 Fascin (Ser-39), phospho-specific Rabbit Polyclonal

AM2021 Actin (C-terminal region) Mouse Monoclonal

AP1651 Actin (N-terminal Region) Rabbit Polyclonal

AP1671 Actin (Tyr-53), phospho-specific Rabbit Polyclonal

FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

web: www.ecmbiosciences.com

telephone: 859-879-2075

email: info@ecmbiosciences.com

toll-free: 1-800-859-8202

ECMBiosciences