

WAVE Phospho-Regulation Antibody Sampler Kit

Catalog # WK6130

Kit Components:

Catalog#	Description	Host	Size	Applications	Species Reactivity	MW (kDa)
WP1771	WAVE (Tyr-125), phospho-specific	Rabbit pAb	50 µl	WB, E	H, R, M	78/80
WP1821	WAVE (Tyr-150), phospho-specific	Rabbit pAb	50 µl	WB, E	H, R, M	78/80
WP1731	WAVE1 (N-terminal Region)	Rabbit pAb	50 µl	WB, E	H, R, M	80
WP1791	WAVE2 (Central Region)	Rabbit pAb	50 µl	WB, E	H, R, M	78
RS3251	Anti-Rabbit Ig Light-Chain Specific:HRP	Mouse mAb	100 µl	WB, E	Rabbit	

Applications: WB = western blot, E = ELISA.

Species: H = Human, R = Rat, M = Mouse, C = Chicken, F = Frog

Kit Summary:

The WAVE phospho-regulation antibody sampler kit can be used to examine phosphorylation of WAVE1 at Tyr-125 and Tyr-150, as well as the conserved sites in WAVE2 and WAVE3. The kit also includes polyclonal antibodies to monitor total expression levels of WAVE1 and WAVE2.

Buffers and Storage:


Rabbit polyclonal antibodies are supplied in 50µl phosphate-buffered saline, 50% glycerol, 1 mg/ml BSA, and 0.05% sodium azide. Store at -20°C. Do not aliquot. Stable for 1 year.

Background:


The Wiskott–Aldrich syndrome protein (WASP) family is involved in various pathways that regulate actin cytoskeletal organization. This family includes WASP, N-WASP, and three WAVE/SCAR isoforms, WAVES 1, 2, and 3. WAVE proteins play key roles in actin-mediated cell events, such as membrane ruffling and lamellipodia formation. WAVES contain an N-terminal WAVE homology domain, a basic domain, a Proline-rich region, and carboxy terminal verprolin, cofilin, and acidic (VCA) region. WAVES are thought to act downstream of the Rac GTPase, connecting Rac activation to induction of Arp 2/3-mediated actin polymerization. Regulation of WAVE activity can occur through tyrosine phosphorylation. Src phosphorylation of WAVE1 at Tyr-125 enhances binding to the Arp2/3 complex, and is required for WAVE inhibition of Arp2/3-mediated stress fiber formation. By contrast, WAVE2 phosphorylation of Tyr-150 by Abl may enhance Arp2/3 complex actin nucleation and microspike formation in fibroblasts. Thus, site-specific tyrosine phosphorylation may be important for controlling specific activities of WAVE proteins.

References:


Ardern, H. et al. (2006) Cell Motil. Cytosk. 63:6.
Leng, Y. et al. (2005) PNAS 102(4):1098
Suetsugu, S. et al. (1999) Bioch. Biophys. Res. Comm. 260:296
Miki, H. et al. (1999) J Biol. Chem. 274(39):27605.



Western blot of human SYF cSrc transformed cells untreated (lanes 1 & 3) or treated (lanes 2 & 4) with pervanadate (1 mM; 30 min). The blots were probed with anti-WAVE1 (N-terminal region) (lanes 1 & 2) or anti-WAVE (Tyr-125) (lanes 3 & 4).



Western blot of human SYF cSrc-transformed cells. Blots were probed with anti-WAVE1 (N-terminal region) at a dilution of 1:1000 (lane 1), 1:2000 (lane 2) or 1:4000 (lane 3). In addition, the antibody was used in the absence (lane 4) or presence of blocking peptides, WAVE1 (N-terminal region) peptide (lane 5) or WAVE2 (Central region) peptide (lane 6).



Western blot of rat PC12 lysate. The blots were probed with anti-WAVE2 (Central region) in the presence (lane 2) or absence (lane 1) of WAVE2 (Central region) blocking peptide (WX1795).

FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

www.ecmbiosciences.com
telephone: 859-879-2075
toll-free: 1-800-859-8202
info@ecmbiosciences.com

ECMBiosciences